

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY) 21/12/2004	2. REPORT TYPE Final Report	3. DATES COVERED July 1, 1998 - June 24, 2002
---	--------------------------------	--

4. TITLE AND SUBTITLE Vascularized Tissue Sensors for Generic Toxin and Pathogen Detection	5a. CONTRACT NUMBER
	5b. GRANT NUMBER N00014-98-1-0760
	5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S) Griffith, L.G., Tannenbaum, S.R., Wands, J., Sherley, J., Schauer, D., So, P.T., Lee, R.T., Kamm, R.D.	5d. PROJECT NUMBER
	5e. TASK NUMBER
	5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Massachusetts Institute of Technology Bldg. 16-429 77 Massachusetts Ave. Cambridge, MA 02139	8. PERFORMING ORGANIZATION REPORT NUMBER
---	---

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5000	10. SPONSOR/MONITOR'S ACRONYM(S) ONR
	11. SPONSOR/MONITOR'S REPORT NUMBER(S)

12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution Unlimited	20050107 075
13. SUPPLEMENTARY NOTES	

14. ABSTRACT
We developed three iterative prototypes of a bioreactor system that facilitates microperfusion of tissue-like structures formed from primary liver cells, and showed that the bioreactor and detection system could be used in a field setting running on batteries. We demonstrated that the endogenous P450 metabolic capacity could be used as a sensor and that cells in the bioreactor were sensitive to toxins requiring liver metabolism.

15. SUBJECT TERMS		
16. SECURITY CLASSIFICATION OF:		
a. REPORT Unclass.	b. ABSTRACT Unclass.	c. THIS PAGE Unclass.
17. LIMITATION OF ABSTRACT UL		
18. NUMBER OF PAGES 5		
19a. NAME OF RESPONSIBLE PERSON Linda G. Griffith		
19b. TELEPHONE NUMBER (Include area code) (617) 253-0013		

FINAL REPORT

GRANT #: N00014-98-1-0760

PRINCIPAL INVESTIGATOR: Linda G. Griffith

INSTITUTION: Massachusetts Institute of Technology

GRANT TITLE: Vascularized Tissue Sensors for Generic Toxin and Pathogen Detection

AWARD PERIOD: 7/1/98 - 6/24/02

OBJECTIVE: To develop an *in vitro* tissue-based sensor based on primary liver cells, and demonstrate the performance of this sensor in detecting a set of toxin including aflatoxin B1, an agent which requires liver metabolism, using a novel optical readout based on cell metabolism.

APPROACH: Building on observations of histotypic 3D reorganization made by previous investigators as well as our own observations regarding reorganization of mixed hepatocyte-endothelial cells cultures, we developed a microfabricated bioreactor that allows us to recreate many features of three dimensional liver tissue at the level of the smallest functional tissue unit, the capillary bed. Further, the design of the reactor allows one to probe tissue structure and function in a noninvasive manner via microscopy. We exploited silicon microfabrication technology to create an array of capillary-bed size channels in a thin silicon chip to serve as a scaffold for 3D tissue morphogenesis. Using a computational fluid dynamics model as a guide, we designed a flow-through reactor housing that allows cells placed in the silicon scaffold to be perfused with nutrient medium at flow rates that provide physiological shear stresses and nutrient delivery rates. The system can be scaled to hold 10^4 - 10^6 cells by adjusting the number of holes in the chip, each of which hold $\sim 10^3$ cells, and by appropriate scaling of the reactor housing. Further, the housing is designed with dimensions that allow the tissue to be within the focal plane of a light microscope or 2-photon microscope so that repeated *in situ* observation of tissue behavior is possible. Test agents can be added to the circulating medium

ACCOMPLISHMENTS

We developed an Integrated sensor that can run on battery power. The final system comprises

(1) a microfabricated bioreactor for differentiated tissue culture with protocols for operation with primary hepatocytes, co-cultures of hepatocytes and endothelial cells, and (co) cultures of liver epithelial cell lines; The reactor is a fourth generation bioreactor (MilliRxF), designed using computational modeling of fluid flow and

mass transfer and incorporating user suggestions from experience with earlier versions.

(2) millifluidic system for distribution of culture medium and reagents with controlled temperature and gas phase humidity/CO₂. This system can be used to culture cells for a period of several days outside the incubator (e.g., room air/temperature).

(3) A miniaturized optical system enabling sequential detection of 3 different fluorescence readouts (EROD/P450; Sytox Nuclear stain; b-lactamase; plus GFP for development work) from one culture within a 10 min measurement period (filter switching required). Four units have been fabricated that are capable of detecting the desired readouts. Results from feasibility studies have led us to focus on P450 (resorufin) as the primary readout for sensor operation, and to GFP for characterization of viral infection.

Performance of the system was demonstrated using aflatoxin, microcystin, shiga toxin

Aflatoxin: Aflatoxin B1 is a natural product of molds that grow on grain, but it is also a potential biowarfare agent. It requires liver metabolism for toxicity, and has a complex metabolic pathway, requiring liver enzymes that are typically rapidly lost in 2D culture. Performance of the system to detect AFB1 toxicity was tested with P450 1A1 ethoxyresorufin readouts. In this assay, the substrate ethoxyresorufin is converted to the fluorescent product resorufin by active intracellular processes, and cell toxicity is reflected by a decrease in signal compared to control. A limit of detection of 5 ng/ml within 24 hours with P450 readout; similar sensitivity was seen for albumin secretion suppression and LDH leakage. An NIEHS grant was obtained to characterize the molecular basis of aflatoxin response in the bioreactor, focusing on transcriptional profiling to compare the bioreactor to the in vivo state in baseline and aflatoxin-treated conditions.

Microcystin: Microcystin is a small peptide toxin produced by cyanobacteria as a natural product, and is found endemic in certain parts of the world in water where bacterial blooms are common. Because microcystin is resistant to filtration and chlorine, it is viewed as a potential biowarfare agent. The system was shown to detect 5 microM microcystin within an hour.

Shiga Toxin: Shiga toxin is a bacterial product that causes loss of blood vessel integrity through action on endothelia. A high degree of variability in setting up endothelial co-cultures limited progress on the Shiga toxin response. Initial co-cultures showed little response by live-dead assay. Further work in assessing toxicity in co-cultures is continuing.

3. Feasibility of Virus detection:

Adenovirus: Adenovirus was chosen as a model virus to characterize the distribution and uptake of a virus in the system. A non-replicating Ad5 vector was used to deliver the gene for eGFP under the CMV promoter, allowing non-invasive analysis of viral uptake and gene expression by measuring fluorescence using the spectrometer. The rate of viral uptake and expression depends on dose, as expected. *This work is being continued under NSF funding.*

CONCLUSIONS: We developed three iterative prototypes of a bioreactor system that facilitates micro-perfusion of tissue-like structures formed from primary liver cells, and showed that the bioreactor and detection system could be used in a field setting running on batteries. We demonstrated that the endogenous p450 metabolic capacity could be used as a sensor and that cells in the bioreactor were sensitive to toxins requiring liver metabolism.

SIGNIFICANCE: Because liver metabolic functions are labile in traditional cultures, the microperfused liver format is a promising advance in providing extended high-level function in culture. The system is being further developed as a means of testing drug toxicity and toxicity of environmental agents.

PATENT INFORMATION:

Griffith, L.G., Tannenbaum, S.R., Powers, M.J., Thompson, C., Domansky, K., "Vascularized Micro Tissue/Micro Organ Arrays," Patent No. 6,197,575

AWARD INFORMATION:

(Griffith, PI)

1999: MIT Class of 1960 Innovation in Education Award

2000: International Fellow, Biomaterials Science and Engineering, International Union of Societies for Biomaterials Science and Engineering

2002: Popular Science Brilliant 10

PUBLICATIONS AND ABSTRACTS.

LINDA G. GRIFFITH

Powers, M.J., and L.G. Griffith "Adhesion-Guided In Vitro Morphogenesis in Pure and Mixed Cell Cultures," Microscopy Res. Technique, 43, 379-384 (1998).

Macdonald, J.M., J.P. Griffin, H. Kubota, L.G. Griffith, J. Fair, L.M. Reid, L.M., Bioartificial Livers, in Cell Encapsulation Technology and Therapeutics, W. Kuhtreiber, R.P. Lanza, and W.L. Chick, eds, Birkhauser, NY (1999).

Griffith, L.G, and A.J. Grodzinsky, "Advances in Biomedical Engineering", JAMA, 285, 556-561 (2001).

Powers, M., Janigian, J., Wack, K.E., Baker, C.E. Stolz, D.B., **Griffith, L.G.**, "Functional behavior of primary rat liver cells in a 3D perfused microarray bioreactor," Tissue Eng., 8, 513-527 (2002).

Powers, M.J., Domansky, K., Kaazempur-Mofrad, M.R., Kalezi, A., Capitano, A., Udupadhaya, A., Kurzawski, P., Wack, K.E., Stolz, D.B., Kamm, R., **Griffith, L.G.**, "A microarray perfusion bioreactor for 3D liver culture," Biotech. Bioeng., 78, 257-269 (2002).

Griffith, L.G. and Naughton, G., "Tissue Engineering: Current Challenges and Expanding Opportunities" Science, 295, 1009-1014 (2002).

Kaazempur-Mofrad, M.R., M. Bathe, H. Karcher, H.F. Younis, H.C. Seong, E.B. Shim, R.C. Chan, D.P. Hinton, A.G. Isasi, A. Upadhyaya, M.J. Powers, L.G. Griffith, and R.D. Kamm, "Role of simulation in understanding biological systems," Computers and Structures, 81, 715-726 (2003).

JAMES L. SHERLEY

Merok, J. L. and Sherley, J. L. (2001) "Breaching the Kinetic Barrier to In Vitro Somatic Stem Cell Propagation," *J. Biomed. Biotech* 1, 24-26.

Rambhatla, L., Bohn, S. A., Stadler, P. B., Boyd, J. T., Coss, R. A., and Sherley, J. L. (2001) "Cellular Senescence: Ex vivo p53-Dependent Asymmetric Cell Kinetics," *J. Biomed. Biotech* 1, 27-36.

Merok, J. R., Lansita, J. A., Tunstead, J. R., and Sherley, J. L. (2002) "Co-segregation of Chromosomes Containing Immortal DNA Strands in Cells That Cycle With Asymmetric Stem Cell Kinetics," *Cancer Research*, 62, 6791-6795.

Lee, H.-S., Crane, G. G., Merok, J. R., Tunstead, J. R., Hatch, N. L., Panchalingam, K., Powers, M. J., Griffith, L. G., and Sherley, J. L. (2003) "Clonal Expansion of Adult Rat Hepatic Stem Cell Lines by Suppression of Asymmetric Cell Kinetics (SACK)", *Biotech. & Bioeng.* 83, 760-771.

Sherley, J. L. (2002) "Asymmetric Cell Kinetics Genes: The Key to Expansion of Adult Stem Cells in Culture", *Stem Cells* 20, 561-572.

Peter T.C. So

Dong, C. Y., French, T., So, P. T. C., Buehler, C., Berland, K. M., Gratton, E., "Fluorescence-lifetime imaging techniques for microscopy," *Methods Cell Biol.*, 72, 431-64 (2003).

Dong, C. Y., Koenig, K., So, P. T. C., "Characterizing point spread functions of two-photon fluorescence microscopy in turbid medium," *J. Biomed. Opt.*, 8, 450-9, (2003).

Ragan, T. M., Huang, H., So, P. T. C., "In vivo and ex vivo tissue applications of two-photon microscopy," *Methods Enzymol.*, **361**, 481-505 (2003).

Dong, C. Y., Buehler, C., So, T. C., French, T., Gratton, E., "Implementation of intensity-modulated laser diodes in time-resolved, pump-probe fluorescence microscopy," *Applied Optics*, **40**, 1109-1115 (2001).

Buehler, C., Dong, C.Y., So, P.T.C., and Gratton, E., "Time-resolved polarization imaging by pump-probe (stimulated emission) fluorescence microscopy," *Biophys. J.*, **79**, 536-549 (2000).

So, P.T.C., "Two-Photon Fluorescence Microscopy: A New Tool for Tissue Imaging and Spectroscopy," *Histotechnology*, **23**, 221-228 (2000).

So, P.T.C., Dong, C.Y., Berland, K.M., and Masters, B.R., "Two-Photon Excitation Microscopy", *Ann. Rev. of Bioeng.*, **2**, 399-429 (2000).

Steven R. Tannenbaum

J. Kim and S.R.Tannenbaum 2003. S-nitrosation regulates the activation of endogenous procaspase-9 in HT-29 human colon carcinoma cells. *Journal of Biological Chemistry* **279** (11), 9758-64

JACK R. WANDS

zu Putlitz J, Tong S, Wands JR. A short region in the genome of hepatitis B virus is critical for maintenance of high transcript levels. *Virology* 1999;254:245-56.

Tong S, Li J, Wands JR. Carboxypeptidase D is an avian hepatitis B virus receptor. *J Virol* 1999;73:8696-702.

Ahn SH, Kramvis A, Kawai S, Spangenberg HC, Li J, Kimbi G, Kew M, Wands JR, Tong S. Sequence variation upstream of precore translation initiation codon reduces hepatitis B virus e antigen production. *Gastroenterology* 2003;125:1370-8.